

Ground-State Properties of Magnetically Trapped Bose-Condensed Rubidium Gas

Gordon Baym¹ and Christopher Pethick^{1,2,3}

¹*Department of Physics, University of Illinois at Urbana-Champaign, 1110 W. Green St.,
Urbana, IL 61801*

²*Nordita, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark,*

³*Institute for Nuclear Theory, University of Washington, Box 351550, Seattle, WA 98195*

Abstract

We give a quantitative account of the ground-state properties of the dilute magnetically trapped ^{87}Rb gas recently cooled and Bose-Einstein condensed at nanokelvin-scale temperatures. Using simple scaling arguments, we show that at large particle number the kinetic energy is a small perturbation, and find a spatial structure of the cloud of atoms and its momentum distribution dependent in an essential way on particle interactions. We also estimate the superfluid coherence length and the critical angular velocity at which vortex lines become energetically favorable.

PACS numbers: 03.75.Fi, 05.30.Jp, 32.80.Pj, 67.90.+z

Typeset using REVTeX

In a remarkable experiment, Anderson et al. [1] have cooled magnetically trapped ^{87}Rb gas to nanokelvin-range temperatures, and observed a rapid narrowing of the velocity distribution and density profile, which is interpreted as the onset of Bose-Einstein condensation. Trapped atom clouds are new systems, beyond liquid ^4He [2] and excitons in semiconductors [3], in which particles obeying Bose-Einstein statistics condense at low temperatures [4,5]; indeed, the condensation of trapped atoms has been a “Holy Grail” of atomic physics [6]. In this paper we give simple quantitative arguments, taking into account effects of the repulsive interatomic interactions, to determine the properties of the quantum ground state of the trapped ^{87}Rb system, including its geometry, momentum distribution, coherence length and critical angular velocity for vortex formation.

In the experiment, the gas is magnetically trapped in an effective three-dimensional harmonic well cylindrically symmetric about the z-axis, with tunable angular frequencies ω_z^0 in the axial (z) direction and $\omega_\perp^0 = \omega_z^0/\sqrt{8}$ in the transverse (xy) plane. The oscillators are characterized by lengths $a_\perp = (\hbar/m\omega_\perp^0)^{1/2}$ and $a_z = (\hbar/m\omega_z^0)^{1/2}$, where m is the atomic mass. During the condensation phase, with $\omega_z^0/2\pi = 208$ Hz, and $a_\perp \approx 1.25 \times 10^{-4}$ cm, the distribution rapidly sharpens with falling temperature, as a macroscopic number, N_0 , of the Rb atoms begin to occupy the lowest mode of the well. In the absence of interparticle interactions the lowest single-particle state has the familiar wave function,

$$\phi_0(\vec{r}) = \frac{1}{\pi^{3/4} a_\perp a_z^{1/2}} e^{-m(\omega_\perp^0 r_\perp^2 + \omega_z^0 z^2)/2\hbar}, \quad (1)$$

where $\vec{r}_\perp$ is the component of $\vec{r}$ in the xy-plane. The density distribution, $\rho(\vec{r}) = N_0 \phi_0(\vec{r})^2$, is Gaussian. However, interatomic interactions strongly modify the particle structure in the well.

The low-energy interactions between polarized ^{87}Rb atoms are repulsive, and are described by an s-wave triplet-spin scattering length, a , determined to be in the range $85a_0 < a < 140a_0$, where a_0 is the Bohr radius [7]. In the limit in which the density varies slowly on a scale a , the interaction energy of the gas per unit volume is given by $E_{int} = (2\pi\hbar^2 a/m)|\rho(\vec{r})|^2$. The repulsive interactions favor a reduction of the density from

the free particle situation, $N_0\phi_0(\vec{r})^2$. As the number of particles increases, the first effect of interactions is to cause the cloud of particles to expand in the transverse direction, where the restoring forces are weaker. With further increase in the number, the cloud expands in the z-direction. The eventual size of the cloud is determined, in the limit in which the inter-particle interactions dominate, by a balance between the harmonic oscillator and interaction energies.

To see the physics of the balance, let us neglect the anisotropy of the oscillator potential and assume that the cloud occupies a region of radius $\sim R$, so that $\rho \sim N/R^3$; then the scale of the harmonic oscillator energy per particle is $\sim m\omega_\perp^2 R^2/2$, while each particle experiences an interaction energy with the other particles $\sim (4\pi\hbar^2 a/m)(N/R^3)$. The characteristic length scale is thus $\sim a_\perp\zeta$, where the dimensionless parameter characterizing the system is

$$\zeta \equiv (8\pi Na/a_\perp)^{1/5} \approx 4.21 \left[(a/100a_0)(N/10^4)(10^{-4}\text{cm}/a_\perp) \right]^{1/5}; \quad (2)$$

under the conditions of the experimental trap with large N , $\zeta \gg 1$. The kinetic energy, on the other hand, is of order $N\hbar^2/(2mR^2)$, so that the ratio of the kinetic to interaction or oscillator energies is of order $\zeta^{-4} \sim N^{-4/5}$; for $N \sim 2000$, the kinetic energy in the condensed phase is a factor ~ 200 smaller than the interaction energies.

To estimate the interaction effects more quantitatively we examine the ground state of the system in terms of its order parameter $\psi(\vec{r})$, where $\int d^3r |\psi(\vec{r})|^2 = N$. [We do not distinguish N and N_0 in the weakly interacting system at zero temperature.] In the Hartree approximation, in which $\psi(\vec{r})/N^{1/2}$ is the lowest single particle mode, the ground state energy of the system is given by a Ginzburg-Pitaevskii-Gross energy functional [8],

$$E(\psi) = \int d^3r \left[\frac{\hbar^2}{2m} |\nabla\psi(\vec{r})|^2 + \frac{m}{2} \left((\omega_\perp^0)^2 r_\perp^2 + (\omega_z^0)^2 z^2 \right) |\psi(\vec{r})|^2 + \frac{2\pi\hbar^2 a}{m} |\psi(\vec{r})|^4 \right]. \quad (3)$$

This approach is familiar in prior studies of Bose-condensed polarized atomic hydrogen [9,10]; see also Refs. [11–13]. The Rb experiments, with lower density, larger atomic mass and stronger interactions, fall, however, in a rather different parameter range.

For a first solution we take ψ in the form of the ground-state wave-function, Eq. (1):

$$\psi(\vec{r}) = \frac{N^{1/2}}{\omega_{\perp}^{1/2} \omega_z^{1/4}} \left(\frac{m}{\pi \hbar} \right)^{3/4} e^{-m(\omega_{\perp} r_{\perp}^2 + \omega_z z^2)/2\hbar}, \quad (4)$$

with effective frequencies, $\omega_{\perp}$ and ω_z , treated as variational parameters. Substitution of (4) into (3) yields the ground state energy

$$E(\omega_{\perp}, \omega_z) = N\hbar \left(\frac{\omega_{\perp}}{2} + \frac{(\omega_{\perp}^0)^2}{2\omega_{\perp}} + \frac{\omega_z}{4} + \frac{(\omega_z^0)^2}{4\omega_z} + \frac{Nam^{1/2}}{(2\pi\hbar)^{1/2}} \omega_{\perp} \omega_z^{1/2} \right); \quad (5)$$

minimizing E with respect to $\omega_{\perp}$ we derive

$$\omega_{\perp} = \omega_{\perp}^0 / \Delta, \quad (6)$$

where

$$\Delta = \left(1 + \frac{\zeta^5}{(32\pi^3)^{1/2}} \left(\frac{\omega_z}{\omega_{\perp}^0} \right)^{1/2} \right)^{1/2}. \quad (7)$$

Interactions, by reducing the effective transverse oscillator frequency by Δ , spread out the distribution in the transverse direction by a factor $\Delta^{1/2}$; when N is sufficiently large that $\zeta \gg 1$, $\Delta^{1/2} \approx 2.55 [(N/10^4)(a/100a_0)(10^4\text{cm}/a_{\perp})]^{1/4} (\omega_z/\omega_{\perp}^0)^{1/8}$.

Spreading in the z-direction begins to become significant when the interaction energy per particle becomes comparable with $\hbar\omega_z$; using (6) and minimizing the resultant ground state energy, $E(\omega_z) = N\hbar (\omega_{\perp}^0 \Delta + \omega_z/4 + (\omega_z^0)^2/(4\omega_z))$, with respect to ω_z , we see that this condition is $(Na/a_{\perp}) \gtrsim (\omega_z^0/\omega_{\perp}^0)^{1/2}$, which is realized under the experimental conditions. In the limit $\zeta \gg 1$, the kinetic energy terms in (5) are negligible, and we find the shift in the frequency in the z-direction,

$$\frac{\omega_z}{\omega_z^0} = \frac{2(\pi\lambda)^{3/5}}{\zeta^2}, \quad (8)$$

where $\lambda \equiv \omega_z^0/\omega_{\perp}^0$; the leading contribution to the energy per particle is

$$\frac{E}{N} = \frac{5\zeta^2}{8\pi^{3/5}} \lambda^{2/5} \hbar\omega_{\perp}^0 \propto N^{2/5}. \quad (9)$$

To obtain the ground state wave function more precisely, we minimize the total energy (3) with respect to ψ , keeping the total number of particles fixed, and thus derive the non-linear Schrödinger equation

$$\left[-\frac{\hbar^2}{2m}\nabla^2 + \frac{1}{2}m(\omega_\perp^0)^2(r_\perp^2 + \lambda^2 z^2) + \frac{4\pi\hbar^2 a}{m}|\psi(\vec{r})|^2 \right] \psi(\vec{r}) = \mu\psi(\vec{r}), \quad (10)$$

where μ is the chemical potential. The physical scales are conveniently brought out by rescaling the lengths, letting $\vec{r} = a_\perp \zeta \vec{r}_1$, and writing $\psi(\vec{r}) = (N/\zeta^3 a_\perp^3)^{1/2} f(\vec{r}_1)$, where $\int d^3 r_1 |f|^2 = 1$; then (10) becomes the dimensionless equation,

$$\left[-\frac{1}{\zeta^4}\nabla_1^2 + r_{1\perp}^2 + \lambda^2 z_1^2 + |f(\vec{r}_1)|^2 \right] f(\vec{r}_1) = \nu^2 f(\vec{r}_1), \quad (11)$$

where $\nu^2 \equiv (2\mu/\zeta^2 \hbar \omega_\perp^0)$.

In the limit of large N , we can obtain an essentially exact expression for the ground state wave function, corresponding to the Thomas-Fermi approximation, by neglecting the kinetic energy term, which falls as ζ^{-4} ; then

$$f(\vec{r}_1)^2 = \nu^2 - r_{1\perp}^2 - (\lambda z_1)^2 \quad (12)$$

in the region where the right side is positive, and $f=0$ outside this region. This form for the wave function is good, except where the density is small, in which case the kinetic energy causes the wave function to vanish smoothly. The normalization condition on f implies that $\nu = (15\lambda/8\pi)^{1/5}$, which translates into the relation between μ and N :

$$\mu = \frac{\hbar\omega_\perp}{2} \left(\frac{15\lambda}{8\pi} \right)^{2/5} \zeta^2 = \frac{\hbar\omega_\perp}{2} \left(\frac{15\lambda N a}{a_\perp} \right)^{2/5}. \quad (13)$$

Since $\mu = dE/dN$, the energy per particle is simply $E/N = (5/7)\mu$, a result smaller than the effective oscillator frequency calculation (9) by a factor $(3600\pi)^{1/5}/7 \approx 0.92$. The central density of the blob is $\rho(0) = m\mu/4\pi\hbar^2 a$.

In the limit of large N , the transverse radius of the cloud is given by $R/a_\perp = (15\lambda N a/a_\perp)^{1/5}$, and the half-height in the z-direction is $Z = \lambda R$. For $N = 2000$, $a = 100a_0$, and $\omega_\perp^0/2\pi = 208/\sqrt{8}$ Hz, one has $R/a_\perp \approx 3.24$. When the oscillator “spring constants” are relaxed in the trap by a factor 75, the equilibrium configuration of the cloud then has $R/a_\perp \approx 2.61$. These increases are consistent with observations [1] made when the cloud is finally released from the trap [14]. We also note that for large N the aspect ratio R/Z equals

λ , whereas in the absence of interactions it is $\lambda^{1/2}$; thus for the experimental conditions, one would expect the aspect ratio to be $\sqrt{8}$.

For the Thomas-Fermi wave function, the momentum distribution is proportional to $|J_2(\kappa)/\kappa^2|^2$, where J_2 is the Bessel function of order 2, $\kappa^2 = (\zeta a_\perp)^2 (p_\perp^2 + (p_z/\lambda)^2)$, and $\vec{p}$ is the particle momentum. When $Na/a_\perp \gg 1$, the width of the momentum distribution is much less than that for a single particle in the oscillator potential.

The sound velocity, c_s , in the interior of the cloud is given by $c_s^2 = (\rho/m)(\partial\mu/\partial\rho) = \mu/m$, which in the large N limit equals $(\hbar\omega_\perp^0)(15\lambda Na/a_\perp)^{2/5}$. In this limit, the lowest mode of excitation in the transverse direction of the system has frequency of order $R/c_s \sim \omega_\perp^0$.

The superfluid coherence length, ξ , which determines the distance over which the condensate wave function can heal, can be estimated by equating the kinetic energy term in Eq. (10), $\sim \hbar^2/(2m\xi^2)$, to the interaction energy, which yields

$$\xi^2 = (8\pi\rho a)^{-1}, \quad (14)$$

where ρ is the local density. With the central density of the cloud computed in the Thomas-Fermi approximation, $\rho(0) = m\mu/(4\pi\hbar^2 a)$, we have

$$\frac{\xi}{R} = \left(\frac{a_\perp}{R}\right)^2 = \left(\frac{a_\perp}{15\lambda Na}\right)^{4/5}. \quad (15)$$

Thus when the number of particles is sufficiently large that the Thomas-Fermi approximation is valid, the coherence length is small compared with the size of the blob, and the system should exhibit superfluid properties more like those of a bulk superfluid than an atomic nucleus, where $\xi \sim R$.

An experimentally important confirmation of Bose-Einstein condensation would be the observation of formation of a vortex line in a rotating system. The critical angular frequency, Ω_{c1} , at which it becomes energetically favorable for a vortex line to be created under rotation about the z -axis is

$$\Omega_{c1} \sim \frac{\hbar}{mR^2} \ln(R/\xi). \quad (16)$$

For cloud radii $\sim 5 \times 10^{-4}$ cm, this value corresponds to a rotation frequency of order 10 Hz.

Finally we consider the case of atoms, such as spin-polarized ^{85}Rb [7] or ^7Li [11], with a negative scattering length, corresponding to a low energy attractive interaction. A uniform state of such atoms at low density would be unstable to formation of long-wavelength density waves, signalling a gas-liquid phase transition. However, as discussed theoretically in [12,13], and seen experimentally in [5], the physics in a trap is different; this can be understood in the present context by considering the variational calculation above. With increasing N , the spatial extent of the wave function is reduced. Provided $\Delta^2 = \left(1 - (2/\pi)^{1/2}(N|a|/a_\perp)(\omega_z/\omega_\perp^0)^{1/2}\right)$ (cf. (7)) remains positive, the kinetic energy term is able to stabilize the system. However, if Δ^2 becomes negative, the attractive forces overwhelm the kinetic energy, and the cloud becomes unstable to collapse. The critical number of particles for collapse is $\sim (\pi/2\lambda)^{1/2}a_\perp/|a|$. In ^{85}Rb , for which $-1000a_0 < a < -60a_0$ [7], under the experimental conditions in Ref. [1] with $\omega_z/2\pi = 208$ Hz, this number is $\sim 20 - 300$; in the ^7Li trap of Ref. [11] it is ~ 3000 . The final state of the collapsed cloud is determined by the shorter-range repulsive components of the interatomic potential.

To summarize, our calculations provide quantitative results that confirm and extend the qualitative considerations in Ref. [1] on the effect of particle interactions on the properties of a cloud of bosons. Experimental confirmation of the dependence on trap parameters, particle number and atomic properties of the size of the cloud and the momentum distribution would give one increased confidence in the interpretation of the data.

We are grateful for the hospitality of the Aspen Center for Physics, where this work was largely carried out. In Aspen we had useful conversations with Andrei Ruckenstein on this topic. We are grateful to Eric Cornell for very helpful instruction on the experiments, and to D. G. Ravenhall for helpful advice. This work was supported in part by NSF Grants NSF PHY94-21309 and NSF AST93-15133, and NASA Grant NAGW-1583 .

REFERENCES

- [1] M.H. Anderson, J. R. Ensher, M.R. Matthews, C.E. Wieman, and E.A. Cornell, *Science* **269**, 198 (1995).
- [2] F. London, *Nature* **141**, 643; *Phys. Rev.* **54**, 947 (1938).
- [3] D.W. Snoke, J.P. Wolfe and A. Mysyrowicz, *Phys. Rev. B* **41**, 11171 (1990); J.-L. Lin and J.P. Wolfe, *Phys. Rev. Lett.* **71**, 1223 (1993); and G.M. Kavoulakis, G. Baym, and J.P. Wolfe, *Phys. Rev. B* (in press).
- [4] A comprehensive review of Bose-Einstein condensation in various systems is given in A. Griffin, D.W. Snoke and S. Stringari, eds, *Bose Einstein Condensation* (Cambridge Univ. Press, Cambridge, 1995).
- [5] C.C. Bradley, C.A. Sackett, J.J. Tollett, and R.G. Hulet, Rice Univ. preprint (1995), have very recently reported evidence of Bose-Einstein condensation in another trapped atomic system, ^7Li .
- [6] K. Burnett, *Science* **269**, 182 (1995).
- [7] J.R. Gardner, R.A. Cline, J.D. Miller, D.J. Heinzen, H.M.J.M. Boesten, and B.J. Verhaar, *Phys. Rev. Lett.* **74**, 3764 (1995).
- [8] V. L. Ginzburg and L. P. Pitaevskii, *Zh. Eksp. Teor Fiz.* **34**, 1240 (1958) *Sov. Phys.-JETP* **7**, 858 (1958); E. P. Gross, *J. Math. Phys.* **4**, 195 (1963).
- [9] V.V. Goldman, I.F. Silvera, and A.J. Leggett, *Phys. Rev. B* **24**, 2870 (1981).
- [10] D.A. Huse and E.D. Siggia, *J. Low Temp. Phys.* **46**, 137 (1982).
- [11] E.R.I. Abraham, W.I. McAlexander, C.A. Sackett, and R.G. Hulet, *Phys. Rev. Lett.* **74**, 1315 (1995).
- [12] P.A. Ruprecht, M.J. Holland, K. Burnett, and M. Edwards, *Phys. Rev. A* **51**, 4704 (1995).

- [13] Y. Kagan, G.V. Shlyapnikov, and J.T.M. Walraven, preprint (1995).
- [14] After submitting this paper, we learned of the complementary work of M. Holland and J. Cooper (preprint), who describe the final “ballistic” expansion of the cloud when it is released from the relaxed trap by numerically solving the time-dependent Gross-Pitaevskii equation, starting from distributions in the relaxed trap corresponding to the present Eqs. (1) and (12).